

Extreme Black Holes: Anabasis and Accidental Symmetry

Achilleas P. Porfyriadis

Harvard University

GR23 Beijing, July 2022



AdS_2 and near-extreme black holes

Near the horizon of (near-)extreme black holes spacetime is AdS_2 -like

Extreme Reissner-Nordstrom; Bertotti-Robinson:

[Bertotti, Robinson (1959)]

$$ds^2 = M^2 \left[-r^2 dt^2 + \frac{dr^2}{r^2} + d\Omega^2 \right], \quad A_t = Mr$$

▶ Applies for a wide class of theories in any D

[Kunduri, Lucietti, Reall (2007)]

- e.g. extreme Kerr in 4D pure Einstein GR

[Bardeen, Horowitz (1999)]

▶ Near-horizon approximations *and* Exact solutions

Anabasis:

Backreaction that destroys the AdS_2 boundary and builds the asymptotically flat region of (near-)extreme BHs.

2012.06562 [JHEP 2103] *with* S. Hadar, A. Lupsasca

“ AdS_2 has no dynamics”

Anti-de Sitter fragmentation

Juan Maldacena, Jeremy Michelson* and Andrew Strominger

*Department of Physics, Harvard University
Cambridge, MA 02138*

*E-mail: malda@bose.harvard.edu, jeremy@bohr.harvard.edu,
andy@planck.harvard.edu*

ABSTRACT: Low-energy, near-horizon scaling limits of black holes which lead to string theory on $AdS_2 \times S^2$ are described. Unlike the higher-dimensional cases, in the simplest approach all finite-energy excitations of $AdS_2 \times S^2$ are suppressed. Surviving zero-energy configurations are described. These can include tree-like structures in which the $AdS_2 \times S^2$ throat branches as the horizon is approached, as well as disconnected $AdS_2 \times S^2$ universes. In principle, the black hole entropy counts the quantum ground states on the moduli space of such configurations. In a nonsupersymmetric context AdS_D for general D can be unstable against instanton-mediated fragmentation into disconnected universes. Several examples are given.

KEYWORDS: Black Holes in String Theory, Conformal Field Models in String Theory, Supersymmetry and Duality.

“AdS₂ has no dynamics”



International Press

*publishers of scholarly mathematical
and scientific journals and books*

Home

Journals

Journal
Content
Online

Books

Information
& Ordering

Company
Contacts

Join Our
Mailing Lists 



Advances in Theoretical and Mathematical Physics

Volume 23 (2019)

Number 2

Rigidity of asymptotically AdS₂ × S² spacetimes

Pages: 403 – 435

DOI: <https://dx.doi.org/10.4310/ATMP.2019.v23.n2.a3>

Authors

Gregory J. Galloway (Department of Mathematics, University of Miami, Coral Gables, Florida, U.S.A.)

Melanie Graf (Faculty of Mathematics, University of Vienna, Austria)

Abstract

The spacetime $AdS_2 \times S^2$ is well known to arise as the ‘near horizon’ geometry of the extremal Reissner–Nordstrom solution, and for that reason it has been studied in connection with the AdS/CFT correspondence. Here we consider asymptotically $AdS_2 \times S^2$ spacetimes that obey the null energy condition (or a certain averaged version thereof). Supporting a conjectural viewpoint of Juan Maldacena, we show that any such spacetime must have a special geometry similar in various respects to $AdS_2 \times S^2$, and under certain circumstances must be isometric to $AdS_2 \times S^2$.

Wider picture on AdS_2 dynamics

- ▶ Backreaction in *asymptotically AdS_2 spacetimes* is problematic.
 - Q: Starting with a linear solution for a scalar ϕ on $AdS_2 \times S^2$, does it extend to a non-linear solution of Einstein-Maxwell-Scalar?
 - A: Not if we insist on an asymptotically AdS_2 solution.
E.g. if we impose Dirichlet boundary conditions on the AdS_2 boundary then backreaction of the scalar on the geometry destroys them.
- ▶ Backreaction in *asymptotically flat spacetimes* makes perfect sense.
 - Q: Starting with a linear solution for a scalar $\phi \sim \sqrt{\epsilon}$ on ERN, does it extend to a non-linear solution of Einstein-Maxwell-Scalar?
 - A: Yes. Generically the fully backreacted nonlinear endpoint is a near-extreme RN with $Q = M\sqrt{1 - \mathcal{O}(\epsilon)}$. [Murata, Reall, Tanahashi (2013)]

The connection of AdS_2 with the asymptotically flat region of BHs allows for consistent backreaction. How? What are the correct boundary conditions?

Perturbations of Bertotti-Robinson

- ▶ Background:

$$ds^2 = M^2 \left[-r^2 dt^2 + \frac{dr^2}{r^2} + d\Omega^2 \right], \quad A_t = Mr$$

- ▶ Spherically symmetric perturbations $(h_{\mu\nu}, a_\mu)$ fully characterized by:

$$h_{\theta\theta} = \Phi_0 + ar + brt + cr \left(t^2 - 1/r^2 \right)$$

Comments:

- ▶ $h_{\theta\theta}$ is gauge invariant under $h_{\mu\nu} \rightarrow h_{\mu\nu} + \mathcal{L}_\xi g_{\mu\nu}$.
- ▶ 4-parameter (Φ_0, a, b, c) family of solutions.
- ▶ Φ_0 parameterizes overall rescaling $M \rightarrow M + \delta M$ with $\Phi_0 = 2M \delta M$.
- ▶ Focus on the remaining triplet:

$$\Phi = ar + brt + cr \left(t^2 - 1/r^2 \right)$$

$SL(2)$ transformation properties

$$\Phi = ar + brt + cr \left(t^2 - 1/r^2 \right)$$

- ▶ The background is invariant under the $SL(2)$ isometries of AdS_2 :

$$H: t \rightarrow t + \alpha$$

$$D: t \rightarrow t/\beta, \quad r \rightarrow \beta r$$

$$K: t \rightarrow \frac{t - \gamma(t^2 - 1/r^2)}{1 - 2\gamma t + \gamma^2(t^2 - 1/r^2)}, \quad r \rightarrow r \left[1 - 2\gamma t + \gamma^2(t^2 - 1/r^2) \right]$$

- ▶ Φ is $SL(2)$ -breaking: (a, b, c) get rotated by the above transformations.
- ▶ However,

$$\mu = b^2 - 4ac \quad \text{is } SL(2)\text{-invariant}$$

- ▶ Using $SL(2)$ transformations one may set

$$\Phi = 2r, \quad \text{when } \mu = 0, \text{sgn}(a + c) = 1$$

$$\Phi = -\sqrt{\mu} rt, \quad \text{when } \mu > 0$$

- ▶ $SL(2)$ -breaking solutions Φ are *not* asymptotically $AdS_2 \times S^2$

Anabasis perturbations

Bertotti-Robinson arises from two physically distinct near-horizon near-extremality scaling limits, $\lambda \rightarrow 0$, of Reissner-Nordstrom

- ▶ Limit #1: Begin with $Q = M$ and put the BH horizon at $r = 0$ (set $M = 1$):

$$ds^2 = - \left(\frac{r}{1 + \lambda r} \right)^2 dt^2 + \left(\frac{r}{1 + \lambda r} \right)^{-2} dr^2 + (1 + \lambda r)^2 d\Omega^2, \quad A_t = \frac{r}{1 + \lambda r}$$

At $\mathcal{O}(1)$ we get Bertotti-Robinson in Poincare coordinates

$$ds^2 = -r^2 dt^2 + \frac{dr^2}{r^2} + d\Omega^2, \quad A_t = r$$

At $\mathcal{O}(\lambda)$ we get, by definition, a linear solution around the above.

$$h_{\theta\theta} = 2r$$

This is the $SL(2)$ -breaking $\mu = 0$ solution $\Phi = 2r$ —Poincare *anabasis solution*

Begins to build the asymptotically flat region of an extreme Reissner-Nordstrom

The nonlinear solution obtained from the $\mu = 0$ perturbation of $AdS_2 \times S^2$, when backreaction is fully taken into account in the Einstein-Maxwell theory, is the extreme Reissner-Nordström black hole.

Anabasis perturbations

- ▶ Limit #2: Begin with $Q = M\sqrt{1 - \lambda^2\kappa^2}$ and put the BH horizon at $\rho = 0$:

$$ds^2 = -\frac{\rho(\rho + 2\kappa + \lambda\kappa\rho)}{(1 + \lambda\kappa)(1 + \lambda\rho)^2}d\tau^2 + \frac{(1 + \lambda\kappa)^3(1 + \lambda\rho)^2}{\rho(\rho + 2\kappa + \lambda\kappa\rho)}d\rho^2 + (1 + \lambda\kappa)^2(1 + \lambda\rho)^2d\Omega^2$$
$$A_\tau = \frac{1}{\lambda} \left(1 - \sqrt{\frac{1 - \lambda\kappa}{1 + \lambda\kappa} \frac{1}{1 + \lambda\rho}} \right)$$

At $\mathcal{O}(1)$ we get Bertotti-Robinson in Rindler coordinates

$$ds^2 = -\rho(\rho + 2\kappa)d\tau^2 + \frac{d\rho^2}{\rho(\rho + 2\kappa)} + d\Omega^2, \quad A_\tau = M(\rho + \kappa)$$

At $\mathcal{O}(\lambda)$ we get, by definition, a linear solution around the above.

$$h_{\theta\theta} = 2(\rho + \kappa)$$

Anabasis perturbations

- ▶ Rindler to Poincare transformation for the Bertotti-Robinson:

$$\begin{aligned}\tau &= -\frac{1}{2\kappa} \ln(t^2 - 1/r^2) \\ \rho &= -\kappa(1 + rt) \\ A &\rightarrow A + d\Lambda, \Lambda = \frac{1}{2} \ln \frac{\rho}{\rho + 2\kappa}\end{aligned}$$

Transforms the Rindler anabasis solution to


$$h_{\theta\theta} = 2(\rho + \kappa) = -2\kappa rt$$

This is the $SL(2)$ -breaking $\sqrt{\mu} = 2\kappa$ solution $\Phi = -2\kappa rt$.

Begins to build the asymptotically flat region of a near-extreme RN

In general, $\Phi = ar + brt + cr(t^2 - 1/r^2)$ with $\mu > 0$, leads to

Rindler anabasis with $\sqrt{\mu} = \sqrt{b^2 - 4ac} = 2\kappa$



The nonlinear solution obtained from the $\mu > 0$ perturbation of $AdS_2 \times S^2$, when backreaction is fully taken into account in the Einstein-Maxwell theory, is the near-extreme Reissner-Nordström black hole with $Q = M\sqrt{1 - \mu/4}$.

Accidental Symmetry:

Coordinate transformation that acts on the perturbative solutions of Einstein equation near extreme black hole horizon

2112.13853 [JHEP 2203] *with* G. Remmen

The linearized Einstein equation

Schematic notation:

- ▶ Background geometry \bar{g} —the Bertotti-Robinson spacetime
- ▶ Metric perturbation h —the Φ solution
- ▶ The linearized Einstein equation as a linear differential operator

$$\mathcal{E}(\bar{g}, h) = 0$$

Consider a finite diffeomorphism

$$(t, r) \rightarrow (t, r) + \lambda \left(\xi^t(t, r), \xi^r(t, r) \right)$$

which transforms both $\bar{g} \rightarrow \bar{g}(\lambda)$ and $h \rightarrow h(\lambda)$.

By general covariance, for *arbitrary* λ and ξ^μ , we have:

$$\mathcal{E}(\bar{g}(\lambda), h(\lambda)) = 0$$

Expanding in λ , we have

$$\mathcal{E}(\bar{g}(0), h(0)) + \lambda \frac{\delta}{\delta \lambda} \mathcal{E}(\bar{g}(\lambda), h(0)) + \lambda \frac{\delta}{\delta \lambda} \mathcal{E}(\bar{g}(0), h(\lambda)) + \mathcal{O}(\lambda^2) = 0$$

Accidental symmetry: definition

Starting with a solution to the linearized Einstein equations around the original background, $\mathcal{E}(\bar{g}(0), h(0)) = 0$, we have

$$\lim_{\lambda \rightarrow 0} [\partial_\lambda \mathcal{E}(\bar{g}(\lambda), h(0)) + \partial_\lambda \mathcal{E}(\bar{g}(0), h(\lambda))] = 0 \quad (1)$$

- ▶ 1st term: hold perturbation fixed, act with a linearized diffeo on the background
- ▶ 2nd term: on fixed background, transform perturbation using linearized diffeo

Equation (1) is valid for any diffeo, i.e. for any ξ^μ .

What if we impose the strong requirement that each term in (1) vanishes individually?

$$\lim_{\lambda \rightarrow 0} \partial_\lambda \mathcal{E}(\bar{g}(0), h(\lambda)) = 0 \quad (2)$$

- ▶ Trivial solutions: Isometries of the background $\bar{g}(\lambda) = \bar{g}(0)$
- ▶ Other solution: *accidental symmetry*—transforms solns h among themselves

Accidental symmetry: electrovacuum case

\mathcal{E} : linearized Einstein-Maxwell equations (electrovacuum)

$\bar{g}(0)$: Bertotti-Robinson

$h(0)$: $\Phi = ar$ ($\mu = 0$ solution)

the solution of $\lim_{\lambda \rightarrow 0} \partial_\lambda \mathcal{E}(\bar{g}(0), h(\lambda)) = 0$ is given by

$$\xi = - \left[\epsilon(t) + \frac{\epsilon''(t)}{2r^2} + \frac{t\epsilon'''(t)}{r^2} \right] \partial_t + \left[r\epsilon'(t) - \frac{\epsilon'''(t)}{2r} \right] \partial_r,$$

where $\epsilon(t)$ is an arbitrary cubic polynomial in t ,

$$\epsilon(t) = e_0 + e_1 t + e_2 t^2 + e_3 t^3.$$

- ▶ $\xi_{0,1,2}$: $SL(2)$ Killing vectors of AdS_2

$$\xi_0 = -(1, 0), \quad \xi_1 = -(t, -r), \quad \xi_2 = -\left(t^2 + \frac{1}{r^2}, -2rt\right)$$

- ▶ ξ_3 : non-trivial accidental symmetry

$$\xi_3 = -\left(t^3 + \frac{9t}{r^2}, \frac{3}{r} - 3rt^2\right)$$

Accidental symmetry: electrovacuum equations

Question: What does ξ_3 do?

Answer: Relates $\mu = 0$ to $\mu \neq 0$. Indeed, we have

$$\Delta\mu = -4a\Delta c = -12\lambda e_3 a^2$$

Accidental symmetries enlarge the possible mappings among solutions to include those beyond the $SL(2)$ isometries, thereby allowing to move from one μ orbit to another.

In spherical symmetry the electrovacuum solutions are constrained by Birkhoff's theorem to the non-propagating degrees of freedom that we have discussed so far.

Can accidental symmetries also turn on propagating d.o.f.?

Accidental symmetry: adding matter

$$\lim_{\lambda \rightarrow 0} \partial_\lambda \mathcal{E}(\bar{g}(0), h(\lambda)) = T \quad (3)$$

Source T must satisfy equations of motion. We consider Klein-Gordon scalar $\square\phi = 0$ s.t. the most general spherically symmetric solution is ($u = t - 1/r$, $v = t + 1/r$)

$$\phi = f_+(v) + f_-(u)$$

Can get solution to (3) from the electrovacuum $\Phi = r$ using the transformation

$$\begin{aligned} \xi^t = & \frac{3}{2r} [F'_+(v) + F'_-(u)] - \frac{3}{2r^2} [F''_+(v) - F''_-(u)] \\ & + \frac{3}{r^3} \left[\int^v \frac{F_+(t_0)}{(t-t_0)^4} dt_0 + \int^u \frac{F_-(t_0)}{(t-t_0)^4} dt_0 \right] \\ & - \frac{1}{r^3} \int^r \int^t \frac{f'_+ \left(\hat{t} + \frac{1}{\hat{r}} \right) f'_- \left(\hat{t} - \frac{1}{\hat{r}} \right)}{\hat{r}} d\hat{t} d\hat{r} \end{aligned}$$

$$\xi^r = r[F'_+(v) - F'_-(u)] - [F''_+(v) + F''_-(u)],$$

where $F_+''''(v) = [f'_+(v)]^2$ and $F_-''''(u) = [f'_-(u)]^2$.

Summary and Remarks

Anabasis: Backreaction that destroys the AdS_2 boundary and builds the asymptotically flat region of (near-)extreme BHs.

- ▶ Q: What is the dual of anabasis in AdS/CFT?

A: Following an inverse RG, from IR to UV, along an irrelevant deformation of the boundary field theory that does *not* respect AdS boundary conditions (e.g. the single-trace $T\bar{T}$ deformation of CFT_2 studied by [Giveon, Itzhaki, Kutasov, et al 2017–])

- ▶ Q: What about JT gravity?

A: $\Phi = \Phi_{JT}$ solves the JT eom $\nabla_\mu \nabla_\nu \Phi_{JT} - g_{\mu\nu} \nabla^2 \Phi_{JT} + g_{\mu\nu} \Phi_{JT} = 0$ on AdS_2 .
 μ = ADM mass of the 2D black holes in JT gravity.

Accidental Symmetry: Coordinate transformation that acts on the perturbative solutions of Einstein equation near extreme black hole horizon and maps them among themselves.

- ▶ Electrovacuum eqs: turn on deviation from extremality
- ▶ Adding KG matter: turn on arbitrary KG source
- ▶ Accidental symmetries may be thought of as “on-shell large diffeomorphisms of AdS_2 ” (made precise in JT gravity)